

NAVAL OCEANOGRAPHIC OFFICE SPECIAL PUBLICATION 1400-NA 2


SURFACE CURRENTS MORWEGIAN AND

Company of the Company

The state of the s

ABSTRACT

THE RESIDENCE OF THE PROPERTY OF THE PROPERTY OF THE PARTY OF THE PART



DISCLAIMER NOTICE

THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

SURFACE CURRENTS

NORWEGIAN AND BARENTS SEAS



JANUARY 1978

REPRINTED 1980

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

NAVAL OCEANOGRAPHIC OFFICE NSTL STATION, MISSISSIPPI 39522

RFACE CURRENTS

WEGIAN AND BARENTS SEAS



JANUARY 1978

REPRINTED 1980

VED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

AVAL OCEANOGRAPHIC OFFICE STL STATION, MISSISSIPPI 39522



ACKNOWLEDGMENTS

Messrs. Raymond J. Beauchesne* and William E. Boisvert made major contributions to this atlas.

^{*}Mr. Beauchesne presently is employed by the Bureau of Naval Personnel.

FOREWORD

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFA IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURF THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UP TIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE PO LATEST COMPUTER TECHNIQUES.

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MOR OF VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON SERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN F OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT A GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR F NAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCRE

> JOHN R. McDONNELL Captain, U.S. Navy

Commander

FOREWORD

S, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, ILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND IES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. O THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERA-VIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE THNIQUES.

ANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA (EARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS /ERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE DEFENSE MAPPING AGENCY, THE DEFENDENT OF THE ATLASES RELY ON THE PERSONAL OBMAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR DEFENDENT OF ALL "WHO IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC ND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

JOHN R. McDONNELL Captain, U.S. Navy Commander ACCESSION for

NTIS White Section TO DDC Buff Section DI UNANNOUNCED DISTRIBUTION/AVAILABILITY CODES

DISTRIBUTION/AVAILABILITY CODES

DIST. AVAIL and/or SPECIAL




SURFACE CURRENT ATLASES

THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD BY ROCKAPPIC FILTER ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WISH MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1403 - 1934. THESE MW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REJION INDEX LIMITS SHOWN BED WEEK, NOO SP 1402-NE TO COVERS MORTH FACIFIC REGION TO EAST OF THE PHILIPPINGS.

THESE SAFET ME ARROL OF THE NIBER OF CHARMS ARE SIR NIP ERRICHIABLE BOSEY OFF


RECENT IMPROVIMENTS IN THE DATA FILE ASSOCIATED INCUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WITE HE MADE



CURRENT ATLASES

AS AMOUNTS OF NEW DATA WARRANT, MOST LIKELY EVERY 12 - 18 MONTHS.

THESE GRAPHICS MAY NOT BE TRULY REPRESENTATIVE OF THE ACTUAL FLOW IN SUCH AREAS AS THE NORTH SEA, PERSIAN GULF, GULF OF THATLAND, AND YELLOW SEA WHERE CURRENTS ARE STRONGLY TIDAL. FOR SUCH AREAS, OTHER SOURCES DESCRIBING PREDICTABLE HOURLY CHANGES OF TIDAL CURRENTS SHOULD BE CONSULTED.



The Surface Current Data File, from which these atlases are derived, consists primarily of over four million ship set and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the United States. The file is supplemented by several thousand Geomagnetic Electrokinetograph (GEK) observations, mostly Japanese. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily United States data.


General Quality

The quality of this data file is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 12 hours) have been eliminated when warranted. Consideration was given to the reliability of the observer; doubtful shipboard computations of set and drift were edited; and observations with erroneous locations (mostly observations on land) have been eliminated. The accepted data are considered most useful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and drift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The drift can be determined along any straight line track and includes all factors which cause changes in the DR position. When a fix is obtained, the current set (direction) is FROM the DR position TO the fix; the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the next observation.


Because the influence of current may vary along a ship's track, the MEAN POSITION of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIF'S DRIFT) OBSERVATION


Data Presentation

The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.






(1) Persistent Current - 60 percent or more of all observations fall within a 45° sector all of the 8-point compass.





(4) <u>Bizonal Flow</u> - Practically all observate concentrated in opposite pairs of sectors, and one pair contains at least opercent as many observations as the pair. This generally indicates variathat occurs in zones of entrainment topposing currents (see examples A and quadrangles 1, 2, and 3).




If there are 12 or more observations in a 1° quadrangle, the surface current is depicted by vector resultants as follows:









- (1) Persistent Current 60 percent or more of all observations fall within a 45° sector of the 8-point compass.
- (2) Prevailing Current 70 percent or more of all observations fall within two adjacent 45° sectors.
- (3) Primary Current with Secondary Direction (a) Primary Current 50 percent or more of all observations fall within three adjacent 45° sectors.
 - (b) Secondary Direction 20 percent or more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 90° of arc.



3% CALMS




- (4) Bizonal Flow Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).
- (5) Variable Current The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.




gle, ere


232









L




· C

ų.







ψį.




بجر



a. i




بكور



74 N	<u></u>	T	T	1	5	Ε		Τ	1	10	D E	Τ	,	, 	15	S E	, 			20) E	,		- -	. DE	j E T	
			ya 1		0 0 300 6																720 q		. 10 2 . 3 6 . 300 9			3 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
									380 0					3 1				3 3	7 a a		3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			9 1		, po 6	•
			0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0				3 6 9 0 140 ¹		2 0 t 0 t	1 0 0 1	370 0	3 9 6 6 9 6 9 6 9 9 9 9 9 9 9 9 9 9 9 9			, o i	4 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	3	5 1 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	10 2 g 5 1 9 37 1	9 0 0	5 3 680 9	2 t	5 1 0 1 2	10 g	\$ 7 1 5 case 0	1 0 10 10 10 10 10 10 10 10 10 10 10 10	#
70 N	200				000 g	1 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 10 1 0 1 0 0 1 0 0 0 0 0 0 0 0 0 0 0					4 0 0		3 0 1 2 04 ²	9 , 6 2 ; 301 ⁰	3 3 1 1 0mc °	31 0 2 gri ⁸	671 8 0 3 671 8	• ;	0 1 0 1	30 C	8 2 1 0 1 0 0			30.0		
70 N				3 t	2 S		3 2 0 0 0 0		9 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	8 2 1 0 2 1 2 2	-	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 1	213 7	12 0 0 30 ⁰	1 3 0 1	3										
		1 J	1 1 0 0 1 0 0 1	6 0 1 0 0 1	one o		3 0 6	7 1 0 10 0	1 0 0 225 1	, , , , , , , , , , , , , , , , , , ,	9 p		0 0 1 100 1	6 3 0 6	3 g	* • • • • • • • • • • • • • • • • • • •											
	0 0 1 2 2 2 2 2 1	9 1 0 1		7 0 0 1 164 ²	5 0 test 1	3 1	0 1 0 1	4 7 9 1 1 3 38 2	,,,	*	1 20 30 1	1 2			·, ;												
	30 0	20 6	7 , 3 m 4	1, 1	3 6 2 cos 9	3 0 0 mm 1	5 9 99	• ;	7 2 0 1 0 0	m.'	1 2 1 20 1	1 0 0 0 2 138 7															
65 N	3 1	3 4	328.6	, i	134.3	•	•,,	12 , 30 °	3 3 135 7	3	= '									·							
64 N O		20 0	, , , en ,	1 a 1	5	E	9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	<u>:</u> ;;;		10	E				15	E	AN	NUA	L-J	20 AN	E THR	oue	H D	EC	25	E	

.



DISTRIBUTION LIST

NAVY

CINCLANTFLT (N37, N353A) COMSECONDFLT COMSIXTHFLT COMNAVICE COMSUBLANT COMNAVAIRLANT COMNAVSURFLANT COMPATWINGSLANT DIRNAVOCEANMET FLENUMWEACEN FLEWEACEN NORFOLK FLEWEACEN ROTA FLEWEAFAC NAVWEASERVFAC JACKSONVILLE NWSED BERMUDA NWSED BRUNSWICK NWSED CECIL FIELD NWSED CHARLESTON NWSED GUANTANAMO BAY NWSED KEFLAVIK NWSED LONDON NWSED MAYPORT NWSED NAPLES NWSED PATUXENT RIVER

NWSED ROOSEVELT ROADS NWSED SIGONELLA NWSED SOUDA BAY

PRIVATE & UNIVERSITIES

FLORIDA ST. UNIV.
LOUISIANA ST. UNIV.
MASS. INST. OF TECH.
ORE. ST. UNIV.
TEXAS AGM UNIV.
UNIV. OF MIAMI
UNIV. OF R.I.
UNIV. OF WASH.
SCRIPPS INST. OF OCEANOGRAPHY
WOODS HOLE OCEANOGRAPHIC INST.

OTHER GOVT.

NOAA/NODC NOAA/NCC

SECURITY CLASSIFICATION OF THIS PAGE (Then Date Entered)	
REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
REPORT NUMBER A A A A	
NOO SPI400 - NA2 / D- /T // X -	8. TYPE OF REPORT & PERIOD COVERED
SURFACE CURRENTS	Final
NORWEGIAN AND BARENTS SEAS	1
	4. PERFORMING ORG. REPORT NUMBER
Au THOR(a)	6 CONTRACT OR GRANT HUMBER(e)
Naval Oceanographic Office	1/14 \ 2 \ 2 \ 7
NSTL Station Ray Sr. Jouig MS 19522 PERFORMING ORGANIZATION NAME AND ADDRESS	(12) 200
	10 PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT HUMBERS
Naval Oceanographic Office (11) 1988	1,
	4
1. CONTROLLING OFFICE NAME AND ADDRESS	January 1978
	13. HUMBER OF PAGES
4. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)	15. SECURITY CLASS. (of this report)
/	
14)/11 M-SP-1400-NA-3	G. DECLASSIFICATION/DOSMORADING
7/11	THE DECLASSIFICATION DOWNGRADING
6. DISTRIBUTION STATEMENT (of this Report)	
7. DISTRIBUTION STATEMENT (of the abotract entered in Black 20, If different in	— Report)
8 SUPPLEMENTARY NOTES	
9. KEY BORDS (Cantinus on reverse olds if necessary and identify by block number	
Surface Currents	
Norwegian Sea Barents Sea	
O. ABSTRACT (Continuo an reverse side if necessary and identify by block mamber)	-
This atlas, and the series of which it is a pand automatically plotted. It makes available to surface current data collected and will be update of data are added to the data file. This and then a wast quantity of data as compared to the predictions printed in the mid-thirties.	o user the most recent and whenever sufficient amounts to other atlases are based
0 1/AH 72 1473 EDITION OF 1 NOV 00 15 OGSOLETE S/H 0102-014-0001 MCCHEVY D.	50450 All

LUSTY CLASSIFICATION OF THIS PAGE (When Date Entered)

20. Continued

The surface current information is based mainly on ship drift, which is the difference between the dead reckoning position and the position determined by any type of navigational fix. This difference describes the direction and speed of the current.

SECURITY CLASSIFICATION OF THIS PAGE/When Date Selected

